

COHEN-MACAULAYNESS OF POWERS OF TWO-DIMENSIONAL SQUAREFREE MONOMIAL IDEALS

NGUYEN CONG MINH AND NGO VIET TRUNG

ABSTRACT. Two-dimensional squarefree monomial ideals can be seen as the Stanley-Reisner ideals of graphs. The main results of this paper are combinatorial characterizations for the Cohen-Macaulayness of ordinary and symbolic powers of such an ideal in terms of the associated graph.

INTRODUCTION

Let I be a two-dimensional squarefree monomial ideal in a polynomial ring $k[X] = k[x_1, \dots, x_n]$, where k is a field and $n \geq 3$. Since we are studying the Cohen-Macaulayness of powers of I , we will assume that I is unmixed. In this case, I is the intersection of prime ideals of the form $P_{ij} := (X \setminus \{x_i, x_j\})$. If we think of the indices of such ideals as the edges of a graph G on the vertex set $\{1, \dots, n\}$, then

$$I = \bigcap_{\{i,j\} \in G} P_{ij}.$$

In other words, I is the Stanley-Reisner ideal of the one-dimensional simplicial complex associated with G [2], [8].

It is well-known that I is a Cohen-Macaulay ideal if and only if G is a connected graph. Our original goal is to characterize the Cohen-Macaulayness of I^m in terms of G for each $m \geq 2$. This kind of problem is usually hard and there are few classes of ideals for which one knows exactly which powers of them are Cohen-Macaulay. Notice that I^m is Cohen-Macaulay for all $m \geq 2$ if and only if I is a complete intersection [1], [10] (A similar characterization for the generalized Cohen-Macaulayness of I^m can be found in [3]). Since the Cohen-Macaulayness of I^m implies the equality $I^{(m)} = I^m$, where $I^{(m)}$ denotes the m -th symbolic power of I , we have to study the problems when $I^{(m)}$ is a Cohen-Macaulay ideal and when $I^{(m)} = I^m$. These problems are of independent interest and they will be solved completely in this paper.

For the Cohen-Macaulayness of $I^{(m)}$ we need to distinguish two cases $m = 2$ and $m \geq 3$. We shall see that $I^{(2)}$ is Cohen-Macaulay if and only if $\text{diam}(G) \leq 2$, where $\text{diam}(G)$ denotes the diameter of G . For $m \geq 3$ we show that $I^{(m)}$ is Cohen-Macaulay if and only if every pair of disjoint edges of G is contained in a cycle of length 4. In that case, all other symbolic powers of I are Cohen-Macaulay, too. The proof is based on a formula for local cohomology modules of quotient rings of monomial ideals found by Takayama [9].

The equality between ordinary and symbolic powers is a purely ideal-theoretic problem. It turns out that there are few graphs which satisfies the condition $I^{(m)} =$

I^m for some $m \geq 2$. In fact, we shall see that $I^{(2)} = I^2$ if and only if $n = 3, 4, 5$ and G is a path or a cycle or the union of two disjoint edges, and for $m \geq 3$, $I^{(m)} = I^m$ if and only if $n = 3, 4$ and G is a path or a cycle or the union of two disjoint edges.

As a consequence of the above results, we show that I^2 is Cohen-Macaulay if and only if $n = 3$ or G is a cycle of length 4, 5, and for $m \geq 3$, I^m is Cohen-Macaulay if and only if $n = 3$ or G is a cycle of length 4. In particular, I must be a complete intersection if I^m is Cohen-Macaulay for some $m \geq 3$.

The results on the equality between ordinary and symbolic powers can be also used to study vertex cover algebras of simplicial complexes recently introduced in [4]. We shall see that the vertex cover algebra of a pure $(n - 3)$ -dimensional simplicial complex Δ on n vertices is standard graded if and only if $n = 3$ or $n = 4$ and Δ is a path or a cycle or the union of two disjoint edges.

1. TAKAYAMA'S FORMULA

Let $S = k[X] = k[x_1, \dots, x_n]$. Let I be an arbitrary monomial ideal in S . Since S/I is an $\mathbb{N}^n$ -graded algebra, $H_m^i(S/I)$ is a $\mathbb{Z}^n$ -graded module over S/I . For every degree $\mathbf{a} \in \mathbb{Z}^n$ we denote by $H_m^i(S/I)_{\mathbf{a}}$ the $\mathbf{a}$ -component of $H_m^i(S/I)$. Inspired of a result of Hochster in the squarefree case [6, Theorem 4.1], Takayama found the following combinatorial formula for $\dim_k H_m^i(S/I)_{\mathbf{a}}$ [9, Theorem 2.2].

For every $\mathbf{a} = (a_1, \dots, a_n) \in \mathbb{Z}^n$ we set $x^{\mathbf{a}} = x_1^{a_1} \cdots x_n^{a_n}$ and $G_{\mathbf{a}} = \{i \mid a_i < 0\}$. We denote by $\Delta_{\mathbf{a}}(I)$ the simplicial complex of all $F \subseteq \{1, \dots, n\}$ such that

- (1) $F \cap G_{\mathbf{a}} = \emptyset$,
- (2) for every minimal generator $x^{\mathbf{b}}$ of I there exists an index $i \notin F \cup G_{\mathbf{a}}$ with $b_i > a_i$.

Let $\Delta(I)$ denote the simplicial complex of all F such that $\prod_{i \in F} x_i \notin \sqrt{I}$.

Theorem 1.1 (Takayama's formula).

$$\dim_k H_m^i(R/I)_{\mathbf{a}} = \begin{cases} \dim_k \tilde{H}_{i-|G_{\mathbf{a}}|-1}(\Delta_{\mathbf{a}}(I), k) & \text{if } G_{\mathbf{a}} \in \Delta(I), \\ 0 & \text{else.} \end{cases}$$

Notice that the original formula in [9] is a bit different. It puts beside $G_{\mathbf{a}} \in \Delta(I)$ the condition $a_i \leq \rho_i - 1$ for $i = 1, \dots, n$, where ρ_i is the maximum of the i th coordinates of all vectors $\mathbf{b} \in \mathbb{N}^n$ such that $x^{\mathbf{b}}$ is a minimal generator of I . But the proof in [9] shows that we may drop this condition, which is more convenient for our investigation.

Using Takayama's formula we obtain the following condition for the vanishing of the first local cohomology module.

Corollary 1.2. $H_m^1(S/I) = 0$ if and only if $\Delta_{\mathbf{a}}(I)$ is connected for all $\mathbf{a} \in \mathbb{N}^n$.

Proof. We need to check when $\tilde{H}_{-|G_{\mathbf{a}}|}(\Delta_{\mathbf{a}}(I), k) = 0$. Therefore, we only need to consider the case $|G_{\mathbf{a}}| = 0$ or, equivalently, $\mathbf{a} \in \mathbb{N}^n$. The conclusion comes from the fact that $\tilde{H}_0(\Delta_{\mathbf{a}}(I), k) = 0$ if and only if $\Delta_{\mathbf{a}}(I)$ is connected. $\square$

As a consequence, if we want to study the Cohen-Macaulayness of a two-dimensional monomial ideal I , we only need to check the connectedness of $\Delta_{\mathbf{a}}(I)$ for $\mathbf{a} \in \mathbb{N}^n$.

We shall see below that there are only a finite number of possibilities for $\Delta_{\mathbf{a}}(I)$. For that we need to reformulate the definition of $\Delta_{\mathbf{a}}(I)$ in a more simple way.

For every subset F of $\{1, \dots, n\}$ let $S_F = S[x_i^{-1} \mid i \in F \cup G_{\mathbf{a}}]$ and $I_F = IS_F$. It is easy to check that $x^{\mathbf{a}} \in S_F$ and $\Delta_{\mathbf{a}}(I)$ is the simplicial complex of all F such that $F \cap G_{\mathbf{a}} = \emptyset$ and $x^{\mathbf{a}} \notin I_F$. If $\mathbf{a} \in \mathbb{N}^n$, we may replace I_F by the ideal obtained from I by setting $x_i = 1$ for $i \in F$.

Using this interpretation of $\Delta_{\mathbf{a}}(I)$ we can show that $\Delta_{\mathbf{a}}(I)$ is closely related to $\Delta(I)$. Notice first that if we denote by P_F the ideal generated by the variables x_i , $i \notin F$, then

$$\sqrt{I} = \bigcap_{F \in \mathcal{F}(\Delta(I))} P_F,$$

where $\mathcal{F}(\Delta(I))$ denotes the set of the facets of $\Delta(I)$.

Lemma 1.3. *$\Delta_{\mathbf{a}}(I)$ is a subcomplex of $\Delta(I)$. Moreover, if I has no embedded associated prime ideals and $\mathbf{a} \in \mathbb{N}^n$, the facets of $\Delta_{\mathbf{a}}(I)$ are facets of $\Delta(I)$.*

Proof. Let F be an arbitrary face of $\Delta_{\mathbf{a}}(I)$. If $F \notin \Delta(I)$, then $\prod_{i \in F} x_i \in \sqrt{I}$. Therefore, $\sqrt{I_F} = S_F$, which contradicts the condition $x^{\mathbf{a}} \notin I_F$. For the second assertion we only need to show that F is contained in a facet of $\Delta(I)$ which belongs to $\Delta_{\mathbf{a}}(I)$. Observe first that $I_F = JS_F$, where J is the intersection of all primary components of I whose associated primes do not contain any variable x_i with $i \in F$. Since $x^{\mathbf{a}} \notin I_F$, there is at least a primary component Q of I such that $x^{\mathbf{a}} \notin Q$. Let G be the set of indices such that P_G is the associated prime of Q . Then $F \subseteq G$. If I has no embedded associated prime ideals, then P_G is a prime component of $\sqrt{I}$, hence G must be a facet of $\Delta(I)$. If $\mathbf{a} \in \mathbb{N}^n$, then $G_{\mathbf{a}} = \emptyset$. Since $x^{\mathbf{a}} \notin QS_G = I_G$, we can conclude that $G \in \Delta_{\mathbf{a}}(I)$. $\square$

Example 1.4. (1) For $\mathbf{a} = 0$ we have $\Delta_{\mathbf{a}}(I) = \Delta(I)$ because for all facets F of $\Delta(I)$ we have $I_F \neq S_F$ and therefore $x^{\mathbf{a}} = 1 \notin I_F$ for all facets F of $\Delta(I)$.

(2) For $\mathbf{a} = \mathbf{e}_i$, where $\mathbf{e}_i$ denotes the i th unit vector of $\mathbb{N}^n$, then the facets of $\Delta_{\mathbf{a}}(I)$ are the facets F of $\Delta(I)$ with the property that x_i is not contained in the primary component of I associated with P_F . In particular, if no primary components of I contains x_i , then $\Delta_{\mathbf{a}}(I) = \Delta(I)$.

The above lemma will play a crucial role in the study of the symbolic powers of two-dimensional squarefree monomial ideals in the next section.

2. COHEN-MACAULAYNESS OF SYMBOLIC POWERS

Let G be a simple graph (i.e. without isolated vertices and loops) on the vertex set $\{1, \dots, n\}$, $n \geq 3$. We associate with G the ideal

$$I_G := \bigcap_{\{i,j\} \in G} P_{ij},$$

where $P_{ij} = (x_t \mid t \neq i, j)$. By definition, the m -th symbolic power of I_G is the ideal

$$I_G^{(m)} := \bigcap_{\{i,j\} \in G} P_{ij}^m.$$

If $n = 3$, I_G is a principal ideal. Hence, $I_G^{(m)} = I_G^m$ and I_G^m is Cohen-Macaulay for all $m \geq 2$. For $n \geq 3$, we will use Takayama's lemma to study the Cohen-Macaulayness of $I_G^{(m)}$.

It is obvious that $\Delta(I_G^{(m)}) = \Delta(I_G)$ for all $m \geq 2$ and that $\Delta(I_G)$ is the one-dimensional simplicial complex associated with G . For simplicity, we identify $\Delta(I_G)$ with G so that the facets of $\Delta(I_G)$ are edges of G . By Lemma 1.3, $\Delta_{\mathbf{a}}(I_G^{(m)})$ is a simple subgraph of G for all $\mathbf{a} \in \mathbb{N}^n$. This graph can be described as follows.

Lemma 2.1. $\Delta_{\mathbf{a}}(I_G^{(m)})$ is the simple graph of the edges $\{i, j\} \in G$ with $\sum_{t \neq i, j} a_t < m$.

Proof. We have to determine which edge $\{i, j\} \in G$ gives a facet of $\Delta_{\mathbf{a}}(I_G^{(m)})$. For $I = I_G^{(m)}$ and $F = \{i, j\}$ we have $I_F = P_{ij}^m$. Since $G_{\mathbf{a}} = \emptyset$ we only need to check when $x^{\mathbf{a}} \notin P_{ij}^m$. But this condition is satisfied if and only if $\sum_{t \neq i, j} a_t < m$. $\square$

Example 2.2. Let $m \geq 2$ and $\mathbf{a} = (m-1)(\mathbf{e}_r + \mathbf{e}_s)$, where $r < s$ are two arbitrary indices. For any edge $\{i, j\} \in G$ we have

$$\sum_{t \neq i, j} a_t = 2(m-1) - a_i - a_j < m$$

if and only if the edge $\{i, j\}$ contains r or s or both. Therefore, $\Delta_{\mathbf{a}}(I_G^{(m)})$ is the subgraph of G which consists of all edges containing r or s or both.

If $I_G^{(m)}$ is Cohen-Macaulay, then $\Delta_{\mathbf{a}}(I_G^{(2)})$ is connected for $\mathbf{a} = (m-1)(\mathbf{e}_r + \mathbf{e}_s)$ by Lemma 1.2. By the description of $\Delta_{\mathbf{a}}(I_G^{(2)})$ in the above example, its connectivity implies that $\{r, s\} \in G$ or there exists an index t such that $\{r, t\}, \{s, t\} \in G$. In other words, the minimal length of paths from r to s is at most 2. Since r, s can be any pair of vertices, G must be connected.

In graph theory, the distance between two vertices of G is the minimal length of paths from one vertex to the other vertex. This length is infinite if there is no paths connecting them. The maximal distance between two vertices of G is called the diameter of G and denoted by $\text{diam}(G)$.

By the above observation, $\text{diam}(G) \leq 2$ if $I_G^{(m)}$ is Cohen-Macaulay. It turns out that this is also a sufficient condition for the Cohen-Macaulayness of $I_G^{(2)}$.

Theorem 2.3. $I_G^{(2)}$ is Cohen-Macaulay if and only if $\text{diam}(G) \leq 2$.

Proof. As we have seen above, we only need to prove the sufficient part. Assume that $\text{diam}(G) \leq 2$. If $I_G^{(2)}$ is not Cohen-Macaulay, there exists $\mathbf{a} \in \mathbb{N}^n$ such that $\Delta_{\mathbf{a}}(I_G^{(2)})$ is not connected by Lemma 1.2. Since $\Delta_{\mathbf{a}}(I_G^{(2)})$ is a simple subgraph of G , $\Delta_{\mathbf{a}}(I_G^{(2)})$ must contain two edges, say $\{1, 2\}$ and $\{3, 4\}$, such that their vertices are not connected by a path. By Lemma 2.1 we have

$$\sum_{t \neq 1, 2} a_t < 2 \quad \text{and} \quad \sum_{t \neq 3, 4} a_t < 2.$$

The first inequality implies that among the components a_t with $t \neq 1, 2$ there is at most a component $a_t \neq 0$ and this component must be 1. Similarly, among the

components a_t with $t \neq 3, 4$ there is at most a component $a_t \neq 0$ and this component must be 1. Therefore, we either have $\mathbf{a} = 0$ or $\mathbf{a} = \mathbf{e}_t$ for some t or $\mathbf{a} = \mathbf{e}_r + \mathbf{e}_s$ with $r \in \{1, 2\}$ and $s \in \{3, 4\}$.

If $\mathbf{a} = 0$ or $\mathbf{a} = \mathbf{e}_t$, then $\Delta_{\mathbf{a}}(I_G^{(2)}) = G$ by Example 1.4, which is a contradiction because G is connected. If $\mathbf{a} = \mathbf{e}_r + \mathbf{e}_s$, then $\Delta_{\mathbf{a}}(I_G^{(2)})$ is the subgraph of G which consists of all edges containing r or s or both. Since $\text{diam}(G) \leq 2$, either $\{r, s\} \in G$ or there exists an index t such that $\{r, t\}, \{s, t\} \in G$. Therefore, $\Delta_{\mathbf{a}}(I_G^{(2)})$ is connected so that we get a contradiction, too. $\square$

For the Cohen-Macaulayness of $I_G^{(m)}$, $m \geq 3$, we need a stronger condition on G .

Theorem 2.4. *For $m \geq 3$, $I_G^{(m)}$ is Cohen-Macaulay if and only if every pair of disjoint edges of G is contained in a cycle of length 4.*

Proof. For simplicity let $I = I_G^{(m)}$. Assume that I is Cohen-Macaulay and that G has two disjoint edges, say $\{1, 2\}$ and $\{3, 4\}$. Consider the vector $\mathbf{a} \in \mathbb{N}^n$ with

$$a_1 = 1, a_2 = 1, a_3 = m - 1, \text{ and } a_i = 0 \text{ for } i = 4, \dots, n.$$

Then $\sum_{t \neq 1, 2} a_t = m - 1$ and $\sum_{t \neq 3, 4} a_t = 2 < m$, hence $\{1, 2\}, \{3, 4\} \in \Delta_{\mathbf{a}}(I)$ by Lemma 2.1. For $i = 4, \dots, n$ we have $\sum_{t \neq 1, i} a_t = m$ and $\sum_{t \neq 2, i} a_t = m$ so that $\{1, i\}, \{2, i\} \notin \Delta_{\mathbf{a}}(I)$. Since $\Delta_{\mathbf{a}}(I)$ is a connected graph by Lemma 1.2, there must be a path in $\Delta_{\mathbf{a}}(I)$ connecting the vertices 1 and 3. This is the case only when $\{1, 3\}$ or $\{2, 3\}$ belongs to $\Delta_{\mathbf{a}}(I)$. From this it follows that $\{1, 3\}$ or $\{2, 3\}$ belongs to G . Similarly, we can show that every vertex of $\{1, 2\}$ or $\{3, 4\}$ is connected at least by an edge of G with the other edge. Now it is easy to see that there are two other edges of G with vertices in $\{1, 2, 3, 4\}$ such that together with $\{1, 2\}, \{3, 4\}$ they form a cycle of length 4.

Conversely, assume that every pair of disjoint edges of G is contained in a cycle of length 4. By Lemma 1.2, if I is not Cohen-Macaulay, there exists $\mathbf{a} \in \mathbb{N}^n$ such that $\Delta_{\mathbf{a}}(I)$ is not connected. Since $\Delta_{\mathbf{a}}(I)$ is a simple graph, $\Delta_{\mathbf{a}}(I)$ must contain two disjoint edges, say $\{1, 2\}, \{3, 4\}$ which belongs to different connected components of $\Delta_{\mathbf{a}}(I)$. By Lemma 2.1, the edges $\{1, 2\}, \{3, 4\}$ belong to G and

$$\sum_{t \neq 1, 2} a_t < m, \quad \sum_{t \neq 3, 4} a_t < m.$$

We may assume that $\{1, 3\}, \{2, 4\} \in G$ in order to get a cycle of length 4 containing the edges $\{1, 2\}, \{3, 4\}$. Since $\{1, 3\}, \{2, 4\} \notin \Delta_{\mathbf{a}}(I)$, we have

$$\sum_{t \neq 1, 3} a_t \geq m, \quad \sum_{t \neq 2, 4} a_t \geq m.$$

The above inequalities yields

$$2m > \sum_{t \neq 1, 2} a_t + \sum_{t \neq 3, 4} a_t = \sum_{t \neq 1, 3} a_t + \sum_{t \neq 2, 4} a_t \geq 2m,$$

a contradiction. So we can conclude that $I_G^{(m)}$ is Cohen-Macaulay. $\square$

Corollary 2.5. *If $I_G^{(m)}$ is Cohen-Macaulay for some $m \geq 3$, then $I_G^{(m)}$ is Cohen-Macaulay for all $m \geq 1$.*

Proof. Since the condition on G of Theorem 2.4 is independent of m , $I_G^{(m)}$ is Cohen-Macaulay for all $m \geq 3$ once $I_G^{(m)}$ is Cohen-Macaulay for some $m \geq 3$. Since this condition implies G is connected with $\text{diam}(G) \leq 2$, it implies that I_G and $I_G^{(2)}$ are Cohen-Macaulay, too. $\square$

Using the above theorems we can easily construct a graph G such that $I_G^{(2)}$ is Cohen-Macaulay but $I_G^{(m)}$ is not Cohen-Macaulay for all $m \geq 3$.

Example 2.6. Let G be a cycle of length 5. Then $\text{diam}(G) = 2$ but no pair of edges of G is contained in a cycle of length 4.

Notice that there are infinitely many graphs which satisfy the conditions of Theorem 2.3 and Theorem 2.4. An instance is the class of complete graphs.

3. EQUALITY BETWEEN ORDINARY AND SYMBOLIC POWERS

Let G be a simple graph on the vertex set $\{1, \dots, n\}$, $n \geq 3$. We want to study the equality between the ordinary and symbolic powers of the ideal

$$I_G := \bigcap_{\{i,j\} \in G} P_{ij},$$

where $P_{ij} = (x_t \mid t \neq i, j)$. Recall that the m -th symbolic power of I_G is the ideal

$$I_G^{(m)} := \bigcap_{\{i,j\} \in G} P_{ij}^m.$$

We observed first that I_G is generated by the monomials $x_i x_j$, where $\{i, j\}$ is a non-edge of G , and $x_i x_j x_k$, where $\{i, j, k\}$ is a triangle of G .

If $n = 3$, then G is either a path or a triangle. Hence I_G is either of the form $(x_1 x_2)$ or $(x_1 x_2 x_3)$, which satisfies the condition $I_G^{(m)} = I_G^m$ for all $m \geq 2$.

If $n \geq 4$, there are only a few graphs which satisfies the condition $I_G^{(m)} = I_G^m$ for some $m \geq 2$. To see that we shall need the following observation.

Lemma 3.1. *Let $n \geq 4$. If G satisfies the condition $I_G^{(m)} = I_G^m$ for some $m \geq 2$, then $n = 4, 5$ and G is a path or a cycle or a union of two disjoint edges.*

Proof. We will first show that if G has a triangle of edges or a triangle of non-edges, then $I_G^{(m)} \neq I_G^m$ for all $m \geq 2$.

If G has a triangle of edges, say with the vertices $\{1, 2, 3\}$, then $x_1 x_2, x_2 x_3, x_1 x_3 \notin I_G$ and $x_1 x_2 x_3 \in I_G$. Since $x_1 x_2 x_3 x_4 \in I_G^{(2)}$, we have

$$x_1^{m-1} x_2^{m-1} x_3^{m-1} x_4 = (x_1 x_2 x_3)^{m-2} x_1 x_2 x_3 x_4 \in I_G^{m-2} I_G^{(2)} \in I_G^{(m)}.$$

On the other hand, if $x_1^{m-1} x_2^{m-1} x_3^{m-1} x_4 \in I_G^m$, then it is divisible by a product of m monomials in I_G . Since only one of them contain x_4 , the others $m - 1$ monomials involve only x_1, x_2, x_3 . Thus, the product of these $m - 1$ monomials is divisible by $(x_1 x_2 x_3)^{m-1}$ so that the remained monomial must be x_4 , a contradiction.

If G has a triangle of non-edges, say with the vertices $\{1, 2, 3\}$, then $x_1 x_2 \in I_G$ and $x_1 x_2 x_3 \in I_G^{(2)}$. Therefore,

$$x_1^{m-1} x_2^{m-1} x_3 = (x_1 x_2)^{m-2} (x_1 x_2 x_3) \in I_G^{m-2} I_G^{(2)} \subseteq I_G^{(m)}.$$

But $x_1^{m-1}x_2^{m-1}x_3 \notin I_G^m$ because it has degree $2m-1$, whereas the minimal degree of the elements of I_G^m is $2m$.

Therefore, if G satisfies the condition $I_G^{(m)} = I_G^m$ for some $m \geq 2$, then G has no triangles of edges and no triangles of non-edges. As a consequence, every vertex belongs to at most two edges and to at most two non-edges of G . For instance, if $\{1, 2\}, \{1, 3\}, \{1, 4\} \in G$, we must have $\{2, 3\}, \{2, 4\}, \{3, 4\} \notin G$, which forms a triangle of non-edges, a contradiction. Using these properties we can easily check that $n \leq 5$ and that G is a path or a cycle in the connected case or the union of two disjoint edges in the unconnected case. $\square$

It turns out that the necessary condition of Lemma 3.1 is also a sufficient condition in the case $m = 2$.

Theorem 3.2. *Let $n \geq 4$. Then $I_G^{(2)} = I_G^2$ if and only if $n = 4, 5$ and G is a path or a cycle or the union of two disjoint edges.*

Proof. By Lemma 3.1, we only need to show the sufficient part. If $n = 4, 5$ and G is a path or a cycle or the union of two disjoint edges, then G has no triangles of edges and no triangles of non-edges. Using this condition we will show that every monomial $f \in I_G^{(2)}$ belongs to I_G^2 .

If f involves only two variables, say x_1, x_2 , then every ideal P_{ij} must contain x_1 or x_2 . Therefore, $\{1, 2\} \notin G$. Hence $x_1x_2 \in I_G$. Since $f \in P_{1j}^2$ for some $j \neq 2$ and since P_{1j} does not contain x_1 , f is divisible by x_2^2 . Similarly, f is divisible by x_1^2 . Hence f is divisible by $(x_1x_2)^2$ so that $f \in I_G^2$.

If f involves only three variables, say x_1, x_2, x_3 , we may assume that $\{1, 2\} \notin G$ and $\{1, 3\} \in G$. Then $x_1x_2 \in I_G$ and $f \in P_{13}^2$. Since P_{13} does not contain x_1, x_3 , f is divisible by x_2^2 . Hence f is divisible by $x_1x_2^2x_3$. If $\{2, 3\} \notin G$, then $x_2x_3 \in I_G$, which implies $f \in I_G^2$. If $\{2, 3\} \in G$, then f is also divisible by x_1^2 . Hence f is divisible by $x_1^2x_2^2$, which implies $f \in I_G^2$.

If f involves more than four variables, we suppose that f is divisible by $x_1x_2x_3x_4$. By the assumption on G we may assume that $\{1, 2\} \in G$ and $\{1, 3\} \notin G$. If $\{2, 4\} \notin G$, then $x_1x_3, x_2x_4 \in I_G$, hence $f \in I_G^2$. If $\{2, 4\} \in G$, then $\{1, 4\} \notin G$. From this it follows that $\{3, 4\} \in G$, hence we must have $\{2, 3\} \notin G$. Therefore, $x_1x_4, x_2x_3 \in I_G$, which implies $f \in I_G^2$. $\square$

For $m \geq 3$, we shall see that the condition $I_G^{(m)} = I_G^m$ implies $n \leq 4$ so that we get the following criterion.

Theorem 3.3. *Let $n \geq 4$. For $m \geq 3$, $I_G^{(m)} = I_G^m$ if and only if $n = 4$ and G is a path or a cycle or the union of two disjoint edges.*

Proof. Assume that $I_G^{(m)} = I_G^m$ for some $m \geq 3$. If $n \geq 5$, then $x_1x_2x_3x_4x_5 \in P_{ij}^3$ for all i, j . Hence $x_1x_2x_3x_4x_5 \in I_G^{(3)}$. We may assume that $\{1, 2\} \notin G$. Then $x_1x_2 \in I_G$. Therefore,

$$x_1^{m-2}x_2^{m-2}x_3x_4x_5 = (x_1x_2)^{m-3}(x_1x_2x_3x_4x_5) \in I^{m-3}I^{(3)} \subseteq I_G^{(m)}.$$

This monomial has degree $2m-1$ so that it can not belong to I_G^m , a contradiction. Thus, $n = 4$. By Lemma 3.1, this implies that G is a path or a cycle or the union of two disjoint edges.

Conversely, if G is one of the graphs in the assertion, then I_G is the edge ideal of the complementary graph of G which consists of the non-edges of G . This graph is a bipartite graph. Hence $I_G^{(m)} = I_G^m$ for all $m \geq 2$ by [7, Theorem 5.9]. $\square$

Now, we can give the following criteria for the Cohen-Macaulayness of I_G^m .

Corollary 3.4. *Let $n \geq 4$. Then I_G^2 is a Cohen-Macaulay ideal if and only if G is a cycle of length 4, 5.*

Proof. This follows from Theorem 2.3 and Theorem 3.2. $\square$

Corollary 3.5. *Let $n \geq 4$. For $m \geq 3$, I_G^m is a Cohen-Macaulay ideal if and only if G is a cycle of length 4.*

Proof. This follows from Theorem 2.4 and Theorem 3.3. $\square$

If $n = 3$ or if G is a cycle of length 4, I_G is a complete intersection. Therefore, I_G^m is a Cohen-Macaulay ideal for some $m \geq 3$ if and only if I_G is a complete intersection. This fact does not hold for $m = 2$ because I_G is not a complete intersection if G is a cycle of length 5.

We can also use the above results to study the vertex cover algebras of certain simplicial complexes.

Let Δ be a simplicial complex on the vertex set $\{1, \dots, n\}$. An integer vector $\mathbf{c} = (c_1, \dots, c_n) \in \mathbb{N}^n$ is called a m -cover of Δ if $\sum_{i \in F} c_i \geq m$ for all facets F of Δ . Let $A_m(\Delta)$ denote the k -vector space generated by all monomials $x^{\mathbf{c}} t^m$ such that $\mathbf{c}$ is a m -cover of Δ , where t is a new variable. Then

$$A(\Delta) := \bigoplus_{m \geq 0} A_m(\Delta),$$

is a graded S -algebra. We call $A(\Delta)$ the vertex cover algebra of Δ [4].

Vertex cover algebra has an interesting algebraic interpretation. Let $\mathcal{F}(\Delta)$ denote the set of the facets of Δ . Then

$$I^*(\Delta) := \bigcap_{F \in \mathcal{F}(\Delta)} P_{\bar{F}},$$

where $\bar{F}$ denotes the complement of F . Then $A(\Delta)$ is the symbolic Rees algebra of $I^*(\Delta)$. It is shown in [4] that $A(\Delta)$ is a finitely generated, graded and normal Cohen-Macaulay S -algebra.

It is of great interest to know when $A(\Delta)$ is a standard graded algebra, that is, when $A(\Delta)$ is generated over S by forms of degree 1. This is equivalent to the condition $I^*(\Delta)^{(m)} = I^*(\Delta)^m$ for all $m \geq 1$ [4] and can be described in terms of the max-flow min-cut property in integer programming [5]. Note that this condition is always satisfied if $\dim \Delta = 0$ ($I^*(\Delta)$ is a principal ideal).

Corollary 3.6. *Let Δ be a pure $(n-3)$ -dimensional simplicial complex on n vertices, $n \geq 4$. Then $A(\Delta)$ is a standard graded algebra if and only if $n = 4$ and Δ is a path or a cycle of length 4 or the union of two disjoint edges.*

Proof. The assumption on Δ implies that the complements of the facets of Δ form a graph G . Since $I^*(\Delta) = I_G$, applying Theorem 3.2 and Theorem 3.3 to G we see that $A(\Delta)$ is a standard graded algebra if and only if $n = 4$ and Δ is a path or a cycle or the union of two disjoint edges. $\square$

REFERENCES

- [1] R. Achilles and W. Vogel, Über vollständige Durchschnitte in lokalen Ringen, Math. Nachr. **1979**, 285-298.
- [2] W. Bruns and J. Herzog, Cohen-Macaulay rings, Cambridge Studies in Advanced Mathematics 39, Cambridge University Press, Cambridge, 1993.
- [3] S. Goto and Y. Takayama, Stanley-Reisner ideals whose powers have finite length cohomologies, Proc. Amer. Math. Soc. **135** (2007), 2355-2364
- [4] J. Herzog, T. Hibi and N.V. Trung, Symbolic powers of monomial ideals and vertex cover algebras, Adv. Math. **210** (2007), no. 1, 304-322.
- [5] J. Herzog, T. Hibi, N.V. Trung and X. Zheng, Standard graded vertex cover algebras, cycles and leaves, Trans. Amer. Math. Soc. **360** (2008) 6231-6249.
- [6] M. Hochster, Cohen-Macaulay rings, combinatorics, and simplicial complexes, in B. R. McDonald and R. A. Morris (eds.), Ring theory II, Lect. Notes in Pure and Appl. Math. **26**, M. Dekker, 1977, 171-223.
- [7] A. Simis, W. Vasconcelos and R. Villarreal, On the ideal theory of graphs. J. Algebra **167** (1994), 389-416.
- [8] R. Stanley, Combinatorics and commutative algebra, Progress in Mathematics 41, Birkhäuser, Boston, 1996.
- [9] Y. Takayama, Combinatorial characterizations of generalized Cohen-Macaulay monomial ideals, Bull. Math. Soc. Sci. Math. Roumanie (N.S.) **48** (2005), 327-344.
- [10] R. Waldi, Vollständige Durchschnitte in Cohen-Macaulay-Ringen, Arch. Math. (Basel) **31** (1978/1979), 439-442.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF EDUCATION, 136 XUAN THUY, HANOI, VIETNAM

E-mail address: ngcminh@gmail.com

INSTITUTE OF MATHEMATICS, 18 HOANG QUOC VIET, HANOI, VIETNAM

E-mail address: nvtrung@math.ac.vn